







where  $M_s$  is the neutron star mass, which is equal to 1.35 Solar masses,  $k_B$  is the Boltzmann constant.

From (5) and (6) it follows that the graviton field cannot heat the center of the star up to its temperature, since part of gravitons passes through the star without transfer of momentum.

The main conclusions obtained in [7] are the following. It is shown that the body mass is defined only by the value of the graviton luminosity, i.e. by the power of energy emission from the body of those gravitons, which interacted with matter and transferred their momentum to it.

The mechanism is presented, by which the magnetars as charged and strongly magnetized neutron stars emit high-energy cosmic rays and neutrinos and photons associated with them. According to the theory of infinite nesting of matter, the analogues of magnetars at the level of elementary particles are protons, which also emit similarly to neutron stars. At the lower level of matter there are praons, that are related to protons just like the latter are related to neutron stars. Praons, protons and magnetars are the densest objects at the corresponding levels of matter, and these objects are the main sources of gravitons, consisting of particles such as neutrinos, photons and cosmic rays. The main contribution into the graviton field, that cause gravitation at the level of stars, is made by the gravitons, produced at the lowest levels of matter. It is assumed that at the level of elementary particles strong gravitation is acting, which is considered as the basis of strong interaction [4]. The value of the strong gravitational constant is many orders of magnitude greater than the ordinary gravitational constant:

$$\Gamma = \frac{e^2}{4\pi\epsilon_0 M_p M_e} = 1.514 \cdot 10^{29} \text{ m}^3 \cdot \text{kg}^{-1} \cdot \text{s}^{-2}, \quad \text{where}$$

$e$  is the elementary charge,  $\epsilon_0$  is the vacuum permittivity,  $M_p$  and  $M_e$  are the masses of the proton and electron, respectively. With the help of the strong gravitational constant we can precisely calculate the radius of the proton and its magnetic moment [9].


Thus it is shown that the graviton field can be the source of gravitation, leading to strengths, scalar and vector field potentials, which are used in the Lorentz-invariant theory of gravitation [4] and in the covariant theory of gravitation [10-11].

### 3 Electromagnetic force


Let us now proceed from gravitation to electromagnetic phenomena. We consider the vacuum field as a multicomponent field, containing neutral neutrinos and photons, as well as energetic charged particles. The field of gravitons is a part of the vacuum field. In order to understand the electric interaction of bodies at a distance from each other, let us consider in Figure 1 the motion of charged small particles in the vicinity of two bodies, one of which is neutral and the second is positively charged.

As can be seen, both positive and negative particles act symmetrically on the positively charged body, which does not result in emerging of any additional force in comparison with the force of gravitation. The same applies to the second neutral body.

Figure 2 a) shows that the positive particles push the negatively charged body to the left, and Figure 2 b) shows that the negative particles push the positively charged body to the right (when the smallest particles pass through the body similarly to gravitons, they transfer their momentum to them). Consequently, both bodies will be attracted to each other.




**Fig. 1.** The lines of motion of the small particles of the vacuum field, which are a) positively charged, b) negatively charged, near two bodies one of which is neutral and the other is positively charged.



**Fig. 2.** The lines of motion of the small particles of the vacuum field which are a) positively charged, b) negatively charged, near two bodies, one of which is negatively charged and the other is positively charged.

Figure 3 shows the lines of motion of the negative particles of the vacuum field near two positively charged bodies. Both bodies attract the negative particles and obtain an additional momentum from them, which leads to repulsion of

bodies. The motion of the positive particles of the vacuum field in Figure 3 is not shown. It is assumed that they are repelled from the bodies and therefore their interaction with them is weak.



**Fig. 3.** The lines of motion of the small particles of the vacuum field, which are negatively charged, near two positively charged bodies.

For two negatively charged bodies the interaction is similar to the one shown in Figure 3, only it is necessary to replace the signs of all charges. This results in the repulsion of similarly charged bodies. The picture described above can be found in [10]. The common in all the Figures is the fact that depending on the sign of the charge of two bodies the number of charged particles falling on the bodies changes so that after calculating the momentum transferred by these particles the electric force in necessary direction arises.

Thus, we reduce the interaction between the charges at a distance to the interaction by means of the charged particles of the vacuum field.

Let us compare the Coulomb's law and the Newton's law:

$$F_C = \frac{q_1 q_2}{4\pi \epsilon_0 R^2}, \quad F_N = \frac{G m_1 m_2}{R^2}.$$

For the charged particles the relation must hold for the fluence attenuation of charged particles in the matter similarly to (2), with substitution of the concentration of nucleons  $n$  with the concentration of the electric charge  $\eta$  inside the body. Instead of the body mass  $m = nM_n$  the absolute value of body charge should be used  $|q| = \eta e$ , where  $e$  is the elementary charge. As a result, instead of (3-4) we arrive at the approximate expression for the vacuum permittivity:

$$\epsilon_0 = \frac{e^2}{6 p_q D_{0q} \mathcal{G}^2} = \frac{e^2}{\epsilon_q \mathcal{G}^2}, \quad (7)$$

where  $p_q$  is the average momentum of a charged particle,  $D_{0q}$  is the fluence rate of the charged particles of the vacuum field,  $\mathcal{G}$  is the cross-section

of interaction of the charged particles with the matter of charged bodies,  $\epsilon_q$  is the energy density of the charged particles in space.

As it was shown in [4], the ratio of the energy density of strong gravitation to the electromagnetic energy density of the proton is equal to the ratio of the proton mass to the electron mass  $\frac{M_p}{M_e}$ . Indeed,

for the field energies and their relations with regard to definition of strong gravitational constant  $\Gamma = \frac{e^2}{4\pi \epsilon_0 M_p M_e}$ , we have:  $E_g = \frac{k \Gamma M_p^2}{R}$ ,

$$E_e = \frac{k e^2}{4\pi \epsilon_0 R}, \quad \frac{E_g}{E_e} = \frac{4\pi \epsilon_0 \Gamma M_p^2}{e^2} = \frac{M_p}{M_e}.$$

We believe that the same relationship holds for the energy density of neutral and charged particles in the vacuum field that allows us to estimate the energy density of charged particles and their cross-section of interaction:

$$\epsilon_q = \epsilon_c \frac{M_e}{M_p} = 4 \cdot 10^{32} \text{ J/m}^3,$$

$$\mathcal{G} = \frac{e}{\sqrt{\epsilon_0 \epsilon_q}} = 2.67 \cdot 10^{-30} \text{ m}^2.$$

This cross-section has the value, which is comparable with the cross-section of nucleon and exceeds the cross-section of gravitons  $\sigma = 5.6 \cdot 10^{-50} \text{ m}^2$ .

If the described picture is true, then from the Coulomb force we can easily move to the field strength of the electric field around a point charge and then to the scalar field potential. After that, dividing the scalar potential by the square of the speed of light and multiplying by the 4-velocity we

obtain the 4-potential of the particle. Then use of the procedure in [11] allows us to find all of the electromagnetic field properties and to derive all the field equations, including the Maxwell equations.

#### 4 Conclusion

After a brief analysis of the models of ether and quantum vacuum and after enumerating the problems existing in these models, we presented the force vacuum field as some alternative. If we assume that the vacuum field consists of such particles as neutrinos, photons and charged high-energy particles, generated at the lowest levels of matter, it helps to explain the high penetrating ability of the particles. The fluxes of gravitons and charged particles of the vacuum field due to the small cross-section of interaction with the matter penetrate all the objects and transfer their momentum to them. Only such dense objects as neutron stars have the ability to appreciably absorb and dissipate the fluxes of gravitons.

According to the estimates in [6], it is necessary to put into a line three neutron stars for significant absorption of the fluxes of gravitons passing through them. Taking into account the fact that the analogues of neutron stars at the atomic level are nucleons and the assumption that the strong gravitation is acting between the nucleons instead of the ordinary gravitation, in [10] and [12] we can explain the effect of saturation of the nuclear forces binding the atomic nuclei. The essence of the explanation lies in the fact that as the number of nucleons increases, the specific energy of strong gravitation per nucleon, which is proportional to the specific nuclear binding energy, stops increasing linearly, as the gravitational field potential usually increases with increasing of the mass. Saturation becomes noticeable in the nuclei, containing about 20 nucleons or more. In these nuclei, due to almost complete absorption of gravitons by nucleons, addition of a new nucleon brings into the system almost the same binding energy, and therefore the dependence of the specific nuclei binding on the atomic number has the saturation effect.

In the presented model the vacuum field is responsible for both gravitational and electromagnetic forces. In contrast to the models of ether and quantum vacuum, in which there is some static substance with certain properties, the vacuum field is a multi-component and dynamic field, consisting of particles moving at about the speed of light. Electromagnetic and gravitational waves in this case must be the waves transferred by the particles of the vacuum field. In particular, in [13]

we have presented the model of a photon, which consists of charged particles.

#### References:

- [1] Dyson F. [Is a Graviton Detectable?](#) Poincare Prize Lecture, International Congress of Mathematical Physics, Aalborg, Denmark, August 6, 2012.
- [2] Caligiuri L.M., Sorli A. Gravity Originates from Variable Energy Density of Quantum Vacuum. *American Journal of Modern Physics*, Vol. 3, No. 3, 2014, pp. 118-128. doi:[10.11648/j.ajmp.20140303.11](#).
- [3] Clark S.J., Tucker R.W. Gauge symmetry and gravito-electromagnetism. *Classical and Quantum Gravity*, Vol. 17, 2000, pp. 4125-4157. doi:[10.1088/0264-9381/17/19/311](#).
- [4] Fedosin S. G. [Fizika i filosofija podobija of preonov do metagalaktik](#). Perm, 1999, 544 pages. ISBN 5-8131-0012-1.
- [5] Fedosin S.G. [Electromagnetic and Gravitational Pictures of the World](#). *Apeiron*, Vol. 14, No. 4, 2007, pp. 385 – 413.
- [6] Fedosin S.G. [Model of Gravitational Interaction in the Concept of Gravitons](#). *Journal of Vectorial Relativity*, Vol. 4, No. 1, 2009, pp. 1 – 24.
- [7] Fedosin S.G. The graviton field as the source of mass and gravitational force in the modernized Le Sage's model. Perm, Preprint, October 2014.
- [8] Fedosin S.G. [The Integral Energy-Momentum 4-Vector and Analysis of 4/3 Problem Based on the Pressure Field and Acceleration Field](#). *American Journal of Modern Physics*. Vol. 3, No. 4, 2014, pp. 152-167. doi: [10.11648/j.ajmp.20140304.12](#).
- [9] Fedosin S.G. [The radius of the proton in the self-consistent model](#). *Hadronic Journal*, Vol. 35, No. 4, 2012, pp. 349 – 363.
- [10] Fedosin S.G. *Fizicheskie teorii i beskonechnaia vlozhennost' materii*. Perm, 2009. ISBN 978-5-9901951-1-0.
- [11] Fedosin S.G. The Procedure of Finding the Stress-Energy Tensor and Equations of Vector Field of Any Form. *Advanced Studies in Theoretical Physics*, Vol. 8, No. 18, 2014, pp. 771 – 779. doi:[10.12988/astp.2014.47101](#).

- [12] Fedosin S.G. [The Principle of Proportionality of Mass and Energy: New Version.](#) *Caspian Journal of Applied Sciences Research*, Vol. 1, No. 13, 2012, pp. 1 – 15.
- [13] Fedosin S.G. [Cosmic Red Shift, Microwave Background, and New Particles.](#) *Galilean Electrodynamics*, Vol. 23, Special Issues No. 1, 2012, pp. 3 – 13.