























The presentation relays more on human verification of the graphs given above. Created algorithm detects climate clusters in the specified time slices and identifies evolution of those clusters over the time. We defined the problem to be solved, provide review of existing solutions and algorithms and analysed whether they could be used in the algorithm created in this paper. Other models such as climate prediction model that apply complex physics calculations use too many correlated parameters and are difficult to derive correct model. The paper gives full procedure of creating an algorithm for solving the defined problem. Creation of the algorithm is given step by step and selections of different mechanisms and/or parameters in that process have been explained.

Also, we presented several ideas for future work in this area, e.g. improving developed solution by implementing mechanisms to improve quality of the results. Combining created algorithm with another could solve more complex problems than the one defined here. For example, implemented algorithm could be combined with another one to identify patterns in climate change in order to predict climate changes. Implemented solution can be used to create data sets for other algorithms (e.g. creating testing data sets for neural network).

#### References:

- [1] David P. Schneider, Clara Deser, John Fasullo and Kevin E. Trenberth, *Climate Data Guide Spurs Discovery and Understanding*, EOS, Transactions American Geophysical Union, 2013, pp. 121-122
- [2] B. Hennemuth, S. Bender, K. Bulow, N. Dreier et al., *Statistical methods for the analysis of simulated and observed climate data, applied in projects and institutions dealing with climate change impact and adaptation*, CSC Report 13, Climate Service Center Germany, 2013
- [3] P. Saikia, H. Tahbilder et al., *Prediction of rainfall using datamining technique over Assam*, Indian Journal of Computer Science and Engineering, 2014
- [4] J. Han and M. Kamber, *Data Mining Concepts and Techniques*, 2nd edition, Morgan Kaufman, 2006
- [5] UN Department of Economic and Social Affairs, Division for Sustainable Development - Area of Climate Change and Sustainable Development, [http://www.un.org/esa/dsd/dsd\\_aofw\\_cc/cc\\_index.shtml](http://www.un.org/esa/dsd/dsd_aofw_cc/cc_index.shtml)
- [6] Nejra Hadzimejlic, Dzenana Donko, Nijaz Hadzimejlic, Climate Data Analysis Using Clustering Data Mining Techniques, *Proceedings of the 3rd International conference on Applied Informatics and Computing Theory (AICT '12)*, WSEAS Press, 2012, pp. 96-100
- [7] P. Kalnis, N. Mamoulis and S. Bakiras, On Discovering Moving Clusters in Spatio-temporal Data, *SSTD proceeding*, 2005, pp. 364-381
- [8] M. Bottcher, F. Hoppner and M. Spiliopoulou, On Exploiting the Power of Time in Data Mining, *ACM SIGKDD Explorations Newsletter 10 (2)*, 2008, pp. 3-11
- [9] S. Günemann, H. Kremer, C. Laufkötter and T. Seidl, Tracing Evolving Subspace Clusters in Temporal Climate Data, *Data Mining and Knowledge Discovery Journal (DMKD)*, Vol. 24, Nr. 2 2012, pp.387-410)
- [10] Klein Tank, A.M.G. and Coauthors, *Daily dataset of 20th-century surface air temperature and precipitation series for the European Climate Assessment*, 2002.
- [11] Project team ECA&D, Royal Netherlands Meteorological Institute KNMI, *Algorithm Theoretical Basis Document v10.4*, European Climate Assessment & Dataset project, 2011
- [12] Matlab R2012a Documentation. Link: <http://www.mathworks.com/help/techdoc/>
- [13] I.H. Witten, E. Frank and M.A. Hall, *Data Mining - Practical Machine Learning Tools and Techniques*, 3rd edition, Morgan Kaufmann, 2011
- [14] C. D. Manning, P. Raghavan and H. Schütze, *Introduction to Information Retrieval*, Cambridge University Press, 2008