

Environmental conditions and oak barrels timber quality: Which is the influence in the *Quercus petraea* Matts. Liebl. forests in the Iberian Peninsula?

PABLO VILA-LAMEIRO, P.; IGNACIO J. DÍAZ-MAROTO

Department of Agroforestry Engineering

High Polytechnic School, University of Santiago de Compostela

Universitary Campus n/n E-27002 Lugo

SPAIN

pablo.vila.lameiro@usc.es

Abstract: - The autoecology of the oak *Quercus petraea* in the northwest Iberian Peninsula was investigated in the present study by applying the methodology developed in previous studies of other species of the genus *Quercus*. For this, the distribution of the species was firstly determined so that a minimum number of representative species could be selected. A total of 52 plots were chosen and in each one, 41 biotype parameters were established (6 physiographic, 17 climatic and 18 edaphic), along with 19 dendrometric and silvicultural parameters. The study of the variability of these parameters allowed description of both central and marginal physiographic-climatic and edaphic habitats of the sessile oak (*Q. petraea*) in the northwest Iberian Peninsula and an assessment of the effect of environmental factors on its present silvicultural status.

The species is indifferent to the type of substrate, tolerates and even prefers lime soils. Within the area of study, sessile oak forests occur at altitudes of between 540 to 1400 m, a range that corresponds to a difference in mean annual temperature of more than 6°C; however, the main climatic feature that defines the stands under study is the abundance of precipitations, with a annual mean precipitation of 1589.8 mm and a summer mean precipitation of 303.6 mm. The species is less resistant to low temperatures than the pedunculate oak (*Q. robur*) and generally shows a longer growth period.

Key-Words: - Environmental / *Quercus petraea* / site / ecological limit / Iberian Peninsula / timber

1. Introduction: The problem

The area of distribution of *Quercus petraea* is very specific and less widespread than *Quercus robur*, being restricted to between latitudes of 37° and 62°N and longitudes of 10° W and 50° E [1, 2, 3], i.e. the easternmost part of Europe [1, 4]. Its distribution is very disperse within the Iberian Peninsula, although wider than that of the pedunculate oak [1, 5, 6], and it is mainly concentrated in the northern mountainous area between Galicia and Cataluña, with notable occurrences in the Cantabrian mountains, the Basque Country and Navarra [4]. In the northwest Iberian Peninsula, it often mixes and hybridizes with *Quercus robur* (*Quercus x rosacea* Bechst) and *Quercus pyrenaica* (*Quercus x*

andegavensis Hy), which makes accurate identification difficult [7, 8, 9].

From a phytosociological point of view, the forests under study are situated in the Eurosiberian Region, Orocantabrian Province, Ubiñense-Picoeuropean and Laciano-Ancarensis sectors, coinciding with the following phytosociological associations [10, 11, 12]: *Linario triornithophorae-Quercetum petraea* (Rivas-Martínez, Izco & Costa ex. F. Navarro 1974) F. Prieto & Vázquez 1987 and *Luzulo henriquesii-Quercetum petraea* (F. Prieto & Vázquez 1987) Díaz & F. Prieto 1994.

The locations where sessile oak forests occur are characterized by abundant precipitation, i.e. more than 600 mm of annual precipitation, of which at least 150 mm falls in the summer [1, 3].

However, the requirements are less stringent than those of *Quercus robur*, allowing the species to colonize drier and wilder sites [13, 14]. The species is not very demanding as regards temperature, tolerating a minimum temperature in the coldest month of around -3°C , and between 15 and 25°C in the hottest one [1, 3]. climatic ranges that are found in Atlantic sites small continental influence [2, 4]. In the northwest Iberian Peninsula, the forests receive an annual precipitation of more than 1100 mm, which may sometimes reach as high as 2000 mm. The summer precipitation is more than 200 mm and in many areas, more than 300 mm [3, 6]. The temperature variations within the study area are low and correspond to a cold temperate climate [15].

The lithologies present in the stands under study are diverse, and develop on both limestone and slates, in soils of high or low fertility, preferentially in loose soils, although they tolerate stony and even rocky soils when the moisture content is high [1, 2, 3]. It's frugalier with soils than *Quercus robur*, and grow mainly on Regosols and Umbrisols [3, 6, 16].

Within the study area *Quercus petraea* occurs in semi-shade and has a fairly robust character [3, 7]. The saplings can grow and develop under cover for several years, but for full development they should be free of cover after about 10 years [13, 17, 18, 19]. They grow slowly, partly because of the low fertility of the soils, at rates ranging between 1 and $5\text{ m}^3\cdot\text{ha}^{-1}\cdot\text{year}^{-1}$ in favourable seasons [3, 6]. Many of these natural forests have been harvested since immemorial time, mainly for naval construction and railway sleepers [1, 3, 5]. In many cases inadequate silvicultural treatments were applied, such as thinning of the best specimens and the unauthorised felling of trees to obtain wood and charcoal [17, 20, 21]. However, because of the difficulties involved in extracting and transporting the wood in mountainous areas, the felling did not have an excessively negative impact, and the forest have suffered more severe deterioration recently, as a result of farming activities and cultivation of the land for agricultural crops [3, 5]. Despite the reduction in the surface area, some highly valuable autochthonous stands have survived, someones in protected sites under the Nature 2000 Network [22]. Another important aspect is the existence of a characteristic flora and fauna, as the sites are the home grounds of species that are in danger of extinction, such as the brown bear and the capercaillie [10, 23], underlining the need to conserve and to try to increase the range of this oak species through sustainable development compatible with conservation.

Background information is provided by the numerous phytosociological studies carried out on these formations [8, 10, 12, 23]. Some authors [3, 4, 6, 7] provide ecological and dendrometric data about *Q. petraea*. The sessile oak is one of the most widely studied broaeaf species within Europe in terms of silviculture and wood quality [17, 19, 20, 24, 25, 26, 27, 28]. In this context, the study of its autoecology, involving analysis of the effect of environmental variables in the development of the species, is fundamental in evaluating the present situation [29, 30, 31]. The aims of the present study were therefore: 1) to characterize the species ecosystems in the northwest Iberian Peninsula, 2) to determine its physiographical-climatic and edaphic habitats, and 3) to relate the environmental variables to silvicultural variables, indicative of its conservation status.

2. Material and methods

2.1. Site description

The study area is located in the northwest Iberian Peninsula, and includes parts of Galicia, Asturias and León (Figure 1). More specifically, the forests under study are mainly located in eastern Galicia, the mountainous area in the south of Asturias and the Leonese side of the Cantabrian Mountain range [3]. comprising the easternmost area of the Orocantabrian Province [8, 11]. The surface area occupied by pure stands of *Quercus petraea* within this territory, estimated on the basis of the most recent data available [31, 32], is approximately 90,000 ha.

2.2. Description of sampling and data recording

Using information provided in the Forest Map of Spain [29] the existing vegetation mosaics in which *Quercus petraea* is present in the area of study were obtained, and the sampling areas were selected from within these, with the help of information provided by Forestry administration personell and data reported in previous studies [3, 6]. To avoid an edge effect, a minimum surface area of one hectare was considered [3, 22, 33]. Prior to the siting of each plot within the stand, the prevailing environmental conditions were established so that a representative site was selected [22]. A total of 52 permanent and rectangular plots (with surface areas of between

400 and 1200 m²) were established on the basis of the stand density (Figure 1 and Table I), so that the number of trees of diameter greater than the minimum that could be inventoried (\emptyset normal ≥ 5 cm) was no less than 50 [34, 35]. The physiographic, dendrometric and soil profile data were then obtained, and along with the climatic data were fitted to the sampling points, according to the methodology of Carballeira et al. [36], and compared with the results of the edaphological analysis, allowing establishment of the biotype parameters in each plot that best define the suitability of the site as a forest site and the present silvicultural status [22, 30, 37].

2.3. Parameters established (see annex)

From a total of 60 parameters studied, 41 were ecological and related to the physiographical structure, climate and edaphology of the biotype and 19 were parameters that characterize *Quercus petraea* forests in terms of dendrometrics and silviculture.

a) *Physiographic parameters* [22, 30, 31, 38]: altitude and mean slope, orientation, sunshine (Gandullo, 1974), depth of soil to the parent rock and distance from the sea.

b) *Climatic parameters* [19, 38, 39, 40].- Precipitation: annual, winter, spring, summer and autumn mean values; Temperature: mean annual temperature, mean temperatures in the hottest and the coldest month, maximum and minimum absolute temperature, temperature range (difference between the maximum t^a and minimum annual) [36], annual evapotranspiration [41] and temperature index [11]; Hydric regime: sum of surpluses and deficits (positive and negative differences respectively, between monthly precipitation and evapotranspiration), hydric index [42] and duration of drought [43].

c) *Edaphic parameters* [22, 29, 30, 31, 44].- Chemical parameters: pH in H₂O, organic matter, nitrogen and ratio of C/N; Parameters that evaluate the soil fertility: available phosphorous, exchangeable potassium, calcium and magnesium. For all of these, the surface concentrations, corresponding to the upper 20 cm of the profile, were determined, except when there was more than one horizon at this depth, in which case the corresponding weighted mean [14, 22, 33] and the total value of the parameter in the profile were calculated, using the method of Russel and Moore [45] (Russell and Moore, 1968). The percentages of fine earth ($\emptyset < 2$ mm) and of total gravel ($\emptyset > 2$

mm) were also calculated.

d) *Dendrometric/silvicultural parameters* [22, 28, 31, 46, 47]: density, number of dead, non-inventoriable and dead specimens of *Q. petraea*, regenerated saplings [3, 14, basimetric area, mean arithmetic and quadratic diameters, standard deviation and coefficient of variation of the height distribution, Assmann's dominant height [35, 48], Hart's index [35, 49] and Czarnowski's index, number of trees in a squared plot of side equal to the arithmetic mean.

2.4. Definition of habitats and statistical analysis

Univariate analysis [50] was applied to the results of all of the parameters under study with the aim of determining the distribution and calculating a series of characteristic values for each parameter [22, 31, 33, 51]: Lower Limit (LL), Lower threshold (LT): 10th percentile, Mean (M), Upper threshold (UT): 90th percentile and Upper limit (UL). The central habitat of *Quercus petraea*, established by recording its presence - and which is comprised between the 10th and 90th percentiles (80% of plots) - allows determination of the most suitable ecological range, and may be considered as the minimum potential area of the species in the study area [22, 31]. Marginal habitats are defined as those between the limits of the central habitat and the absolute extremes; they include 20% of the remaining plots and provide an estimation for the species, of the parameters that remain outside of the habitat, as not all have the same level of significance as biotype descriptors [22, 52, 53]. It was therefore necessary to carry out discriminant analysis of the plots to identify those parameters that have a greater descriptive weight [13, 54, 55, 56], using the methodology proposed by Hill [57, 58], which allows dichotomous classification of the randomly sampled plots [59], such as the presence of plant species [33, 54, 60] and/or the values of certain parameters [14, 22, 29, 53], using Community Analysis Package (CAP) software, more specifically the Twinspan programme [58, 61].

Initial bivariate analysis was carried out using the silvicultural parameters [62] to select which define the present status of the *Quercus petraea* forests in the northwest Iberian Peninsula. The groups classified using Twinspan were then ratified by considering the selected silvicultural and ecological parameters together or using only those obtained in the discriminant analysis [22, 31]. Principal Components Analysis (PCA) was then applied using the method of extraction by squared

minima, creating new variables by combining parameters, which were assigned a value for each plot and were represented spatially, comparing them with the groups created by Twinspan [31, 62, 63].

3. Results and Problem solution

The physiographic data corresponding to the sampled plots are shown in Table 1. In most cases, the stands occur at altitudes above 1000m and with steep slopes, sometimes extremely steep (>75%), and predominantly in the shade.

To characterize the habitat of the oak *Q. petraea* on a regional scale in the northwest Iberian Peninsula, the values corresponding to the parameters selected in the study of biotype and silvicultural status were considered as reference data. The univariate analysis of the physiographic and climatic descriptive statistics (Table II) revealed that the highest variability was obtained in the ORI and CMT parameters, with coefficients of variation (CV) higher than 70% and, in some extreme cases, i.e. AmT and DD, the values were higher than 100 and 200%, respectively. The lowest variability corresponded to TR, AE and HI, with coefficients lower than 10%. Of the 18 edaphic parameters, 12 showed coefficients of variation higher than 40% and in some of these (sP, Ca and sCa), higher than 100%; only the total and surface pH total showed low variability, with coefficients of variation lower than 10% (Table III). Within the silvicultural statistics, the high CV of the parameters NNQ, NDQ, TNNT, TNNT and REG were notable. In contrast, the height variables (MAH, MQH and ADH), the dominant diameter and the basimetric area showed the lowest variability (Table IV).

The physiographic, climatic and edaphic habitats in both central and marginal areas (Figures 2 and 3) were then determined, and in order to identify the most discriminatory ecological parameters in the habitat of the sessile oak, a dichotomous classification of the inventoried plots was made by applying the CAP-Twinspan procedure, using all of the ecological parameters considered (in total 41), because any of these may be discriminatory within the area occupied by the species in the northwest Iberian Peninsula. As a result of the classification, 12 groups of plots were obtained (Table V), which were characterized by physiographic (ALT), climatic (AP and TI), and above all, edaphic parameters (spH, sK, sOM and Ca). Bivariate analysis was then used to select the silvicultural parameters that showed the most significant interrelationships, i.e. NT/h, MAD, MQD, MAH, ADH, MQH and DOD (Table VI). In

theory these variables should be the best descriptors of use and present status of *Q. petraea* forests in the area of study.

Finally, PCA was used to create five new variables from combinations of the previous parameters, which account for more than 79% of the existing variability and the significance of which can be established by with the maturity of the stand: surface pH, surface potassium, altitude of the stand and continentality of the site. By assigning the value of these variables to each plot and creating a spatial representation of the most significant (maturity, spH and sK), was obtained the distribution of the groups resulting from the Twinspan classification (Fig.4).

4. Discussion and conclusions

4.1. Univariate analysis of the physiographic and climatic parameters

The lack of drought in most of the plots gave rise to extreme variation in the duration of drought (DD) parameter, ranging between 0 and 1 in all plots (Table II) [31, 36, 39]. The pattern in the values of mean annual temperature of the absolute minima (AmT) and of the mean temperature in the coldest month (CMT), was previously observed in a study of the autoecology of *Quercus robur* in Galicia [22], and the wide temperature range is accentuated by the range of altitudes within the study area [3, 6]. Despite the existing variation in the orientation, shady locations predominate (Table I), independently of the robust nature of the species [1, 3, 7], as many of these stands have survived in sites where felling would be difficult because of the physiographic conditions [3, 6, 10, 23]. In contrast, the parameters TR, AE and HI show a CV < 10% and that corresponding to the remaining climatic parameters was rarely higher than 30%, which may indicate the existence of a fairly homogenous climate of humid oceanic type, with some continental influence in certain areas [3, 8, 12, 15]

The high variability in several edaphic parameters (Table III), even more evident than in *Quercus robur* [14, 22], is due to the wide range of substrates existing, with lithologies of siliceous nature and limestone, although the pH values show little variation, corresponding to acidic soils [64] (Wilde, 1946).

4.2. Physiographic-climatic habitat of *Quercus petraea* in the northwest Iberian Peninsula

As regards the central habitat (Figure 3), the stands of sessile oak are located in areas of high altitude and slope, in which the orography of the land makes felling of the trees difficult and favours their conservation [3, 6]. Northern orientations predominate, in accordance with the values for the sunshine parameter [65]. The soil may be considered deep, except in the areas of steepest slope. The continentality is not very apparent and is compensated for by the high altitudes [8, 36]. The central values of the annual precipitation range between 1200 and 1850 mm, with precipitations in summer between 150 and 227 mm, indicative of a high demand for moisture, although somewhat less than in *Quercus robur* [14, 22], as demonstrated by the values of the hydric index, which indicate a perhumid climate [42] and by the duration of drought, which is non-existent in most of the stands [3, 6, 8]. The central habitat varies widely in terms of temperature parameters (Figure 3), and thus the temperatures in the study area appear to be ideal for *Quercus petraea* [22, 51] and allow the climate to be classified as mesothermic [41] according to the values of ETP.

Analysis of the marginal habitats (Figure 3) shows that certain parameters are not suitable for fixing the lower value of the central habitat, given the extent of the lower marginal intervals [22, 31, 33], specifically the temperature index (TI) and the sum of the surpluses (SUR). For the remaining parameters there were few plots outwith the central habitat.

4.3. Edaphic habitat of *Quercus petraea* in the northwest Iberian Peninsula

The edaphic parameters surface phosphorous and total potassium, as well as both total and surface calcium, (Figure 4), did not allow accurate determination of the upper limit of the central edaphic habitat of the sessile oak, due to the wide range of the upper marginal intervals [14, 29, 31]. Furthermore, the central values of the parameters C/N, sC/N and P ranged widely, which may indicate that edaphic conditions are very suitable for the development of the species [3, 6, 22]. In the remaining parameters, the most relevant characteristics are as follows (Figure 4):

- The soils are developed on different lithologies, both siliceous and limestone, however, in general they are strongly acidic [64]

(Wilde, 1946) with a little variation of the pH values [3, 6, 30]. According to the FAO classification [16], they belong to five different groups, with a clear dominance of Umbric Regosols, which represent almost 79% of all of the soils studied. Humic Umbrisols account for 12% of the total and finally, Cambisols, Dystric Regosols and Lithic Leptosols each have a single representative. This shows that most of the stands inventoried grow on less well developed soils than *Quercus robur* forests [14, 22].

- There is little variation between the total and surface values of the percentage of organic matter, and the sOM was higher than 20% in only a few plots, which in combination with the existing granulometric composition gives rise to permeable soils [3, 5].

- Although the mean value of C/N ratio was close to 18, the low values of pH do not allow optimal moisture conditions, giving rise to a moderate type humus in most of the plots [8, 30, 37].

- The concentrations of macronutrients, except for phosphorous, show higher values than those observed in *Quercus robur* forests, indicating intermediate soil fertility [14, 22, 66]. The concentrations of phosphorous were also lower than obtained by other authors for these stands [66, 67], which may indicate that the areas presently occupied by sessile oak forests have not undergone changes in land use [3, 6] because this macronutrient is highly associated with rocks containing apatite minerals, which are scarce in the study area.

4.4. Silvicultural characteristics of *Quercus petraea* forests

The stands under study show a wide range of ages and qualities as a result of the different harvesting techniques to which they have been subjected, and at present pure stands do not exist [3, 6, 14]. The descriptive statistics corresponding to the 19 dendrometric/silvicultural parameters used (Table IV) provide an idea of the heterogeneity existing, mainly in the regenerated, non-inventoriable and dead trees, with coefficients of variation that are much higher than the mean value. Although the presence of non-inventoriable specimens of sessile oak is rare, the mortality is high, in contrast with the accompanying species, which adapt well to the closed undergrowth with little available light in these forests - at least in the early stages of development [2, 3, 17, 18, 40]. The coefficients of variation for the remaining parameters, except for density and Czarnowski's index, were lower than 40% (Table IV), with the least variation

corresponding to the parameters MQH, MAH, DOD, BA and ADH [17, 22, 25].

Most of the stands sampled are located in low-lying areas, are well adapted to the surroundings and lack health problems, but have not been managed, because of the lack of any tradition of carrying out silvicultural treatments on slow growing broadleaf species in the northwest Iberian Peninsula [3, 6]; it would therefore be necessary to convert them into regular or semirregular forest before undertaking their management [14, 22], however in this case, management would be more complex [17, 20, 40]. The development of productive silviculture for *Quercus petraea* in the study area may only be considered in areas in which the site quality is intermediate-high [3].

4.5. Relationship between silvicultural parameters and biotype

The dendrometric/silvicultural parameters that would in theory be most suitable for comparison with the biotype parameters are those that were significantly related to all of the other parameters [22, 38, 44], in this case, NT/h, MAD, MQD, MAH, ADH, MQH and DOD (Table VI). Of these, the coefficient of variation for the number of trees per hectare (NT/h) was higher than that corresponding to others (Table IV) and the dominant height (ADH) was affected by the silvicultural treatments carried out, some with negative effects on the correct development of the trees, such as crown cropping [22], and thus it was considered not appropriate to use these in the factorial analysis comparing with the ecological parameters, and only the discriminant functions obtained by applying the Twinspan procedure were used (ALT, PI, IT, PHS, KS, MOS and Ca) (Table V).

The principal components analysis identified stand maturity, spH and sK as the factors that best explain the variability in the stands analyzed [3]. The results obtained, in relation to the grouping of plots, derived from the PCA (Figure 4) are consistent with the dichotomous classification obtained with the Twinspan procedure, with the repetition of groups such as “H” and “I” (Table V) (plots in the Asturian mountains), “E” (plots in Os Ancares in Galicia and Leon) and “L” (plots in Leonese valleys) being notable. However, the groups D”, “F” and “K” appear disperse, because although they are geographically very similar, as reflected by the Twinspan analysis, the stands are very different.

Acknowledgements

The present study was financed by the collaboration between Pernod Ricard and Irish Distillers Inc. with the research group GI-1720 of the Department of Agroforestry Engineering of the University of Santiago de Compostela.

References:

- [1] Ceballos L., Ruiz de la Torre J., 1979. Árboles y arbustos, E.T.S.I. Montes, Madrid.
- [2] Bary-Lenger A., Nebout J.P., 1993 Le chêne pédonculé et sessile en France et en Belgique, Gerfaut Club, Editions du Perron, Allier-Liège.
- [3] Vila P., 2003. Estudio epidométrico y xilológico de las masas de *Quercus petraea* (Mattuschka) Liebl. en el noroeste de la Península Ibérica, Tesis Doctoral, Universidade de Santiago de Compostela.
- [4] Amaral J., 1990. *Quercus*, En: Castroviejo, S. et al., (eds.), Flora Ibérica II, C.S.I.C., Madrid.
- [5] Ruiz de la Torre J., 1991. Mapa Forestal de España, Dirección General de Conservación de la Naturaleza, Instituto Geográfico Nacional, Ministerio de Medio Ambiente, Madrid.
- [6] Vila P., Díaz-Maroto I.J., 2002. Las masas actuales de *Quercus petraea* en Galicia, Inv. Agr. Sis. Rec. For. Vol. 11(1): 5-29.
- [7] Silva-Pando F.J., Rigueiro A., 1992, Guía das árbores e bosques de Galicia, Galaxia.
- [8] Amigo J., Romero M.I., 1994. Vegetación atlántica bajo clima mediterráneo: un caso en el noroeste ibérico, Phytocoenología 22 (4): 583-603.
- [9] Kremer A., Dupouey J.L., Deans J.D., Cottrell J., Csaikl U., Finkeldey R., Espinel S., Jensen J., Kleinschmit J., Van Dam B., Ducouso A., Forrest I., de Heredia U.L., Lowe A.J., Tutkova M., Munro R.C., Steinhoff, S. Badaeu V., 2002. Leaf morphological differentiation between *Quercus robur* and *Quercus petraea* is stable across western European mixed oak stands, Ann. Sci. For. 59 (7):

777-787.

[10] Fernández Prieto J.A., Vázquez V., 1985. Datos sobre los bosques asturianos orocantábricos occidentales, *Lazaroa* 7: 363-382.

[11] Rivas-Martínez S., 1987. Memoria y mapas de series de vegetación de España 1:400.000, ICONA, Ministerio de Agricultura, Pesca y Alimentación, Madrid.

[12] Díaz T.E., Fernández Prieto J.A., 1994. La vegetación de Asturias, *Itinera Geobotánica* 8: 243-528.

[13] Timbal J., Aussenac G., 1996. An overview of ecology and silviculture of indigenous oaks in France, *Ann. Sci. For.* 53 (2-3): 649-661.

[14] Díaz-Maroto I.J., 1997. Estudio ecológico y dasométrico de las masas de carballo (*Quercus robur* L.) en Galicia, Tesis doctoral, Universidad Politécnica de Madrid.

[15] Allué-Andrade J.L., 1990. Atlas Fitoclimático de España. Taxonomías. Ministerio de Agricultura, Pesca y Alimentación, INIA, Madrid.

[16] FAO, 1999. World Reference Base for Soil Resources, *World Soil Resources Reports*, 84.

[17] Jarret P., 1996. Sylviculture de chêne sessile, *Bulletin technique* 31: 21-29, Office National des Forêts.

[18] Kelly D.L., 2002. The regeneration of *Quercus petraea* (sessile oak) in southwest Ireland: a 25-year experimental study, *For. Ecol. Manag.* 166: 207-226.

[19] Lebourgeois F., Cousseau G., Ducos Y., 2004. Climate-tree-growth relationships of *Quercus petraea* Mill. stand in the forest of Bercé ("Futaie des Clos", Sarthe, France), *Ann. Sci. For.* 61(4): 361-372.

[20] Le Goff N., 1984. Indice de productivité des taillis-sous-futaie de chêne dans la région Centre, *Ann. Sci. For.* 41(1): 1-34.

[21] Hochbichler E., 1993. Methods of oaks silviculture in Austria, *Ann. Sci. For.* 50 (6): 583-591.

[22] Díaz-Maroto I.J., Vila P., Silva-Pando F.J., 2005. Autoecology of oaks (*Quercus robur* L.) in Galicia (Spain), *Ann. Sci. For.* 62 (7) (in press).

[23] Fernández Prieto J.A., Bueno A., 1996. La reserva integral de Muniellos: flora y vegetación, Consejería de Agricultura, Principado de Asturias.

[24] Zhang S.Y., Nepveu G., Eyono R., 1994. Intratree and intertree variation in selected wood quality characteristics of European oaks (*Quercus petraea* and *Quercus robur*), *Can. J. For. Res.* 24: 1818-1823.

[25] Duplant P., 1997. Croissance en hauteur dominante du chêne sessile (*Quercus petraea* Liebl.) en futaie régulière, *Bulletin technique* 33: 49-58, Office National des Forêts.

[26] Bergès L., Hervé J.C., Franc A., Gilbert J.M., Nepveu G., 1999. Influence of ecological factors and individual effects on radial growth and wood density components for sessile oak (*Quercus petraea* Liebl.) in Paris Basin and North-Eastern by use of mixed models, *Proceedings Workshop IUFRO S5.01-04*: 205-222.

[27] Bèrges L., 2000. Variabilité individuelle et collective de la croissance et de la densité du bois de *Quercus petraea* en relation avec les facteurs écologiques, Thèse doctorale, École National des Eaux et Forêts, INRA, Nancy.

[28] Polge H., 1973. Facteurs écologiques et qualité du bois, *Ann. Sci. For.* 30(3): 307-328.

[29] Rubio A., Escudero A., Gandullo J.M., 1997. Sweet chestnut silviculture in a ecological extreme of its range in the west of Spain (Extremadura), *Ann. Sci. For.* 54(7): 667-680.

[30] Gandullo J.M., Sánchez O., González S., 1983.

- Estudio ecológico de las tierras altas de Asturias y Cantabria, Monografías INIA 49, Madrid.
- [31] Blanco A., Rubio A., Sánchez O., Elena R., Gómez V., Graña D., 2000. Autoecología de los castaños de Galicia (España), *Inv. Agr. Sis. Rec. For.*, Vol. 9 (2): 337-361.
- [32] DGCONA, 2003. Tercer Inventario Forestal Nacional, Principado de Asturias, Ministerio de Medio Ambiente.
- [33] Gandullo J.M., Bañares A., Blanco A., Castroviejo M., Fernández A., Muñoz L., Sánchez O., Serrada R., 1991. Estudio ecológico de la laurisilva canaria, ICONA.
- [34] Hummel F.C., 1959. Code of Sample Plot Procedure, Forestry Commission Booklet 34.
- [35] Rondeux J., 1993. La mesure des arbres et des peuplements forestiers, Les Presses Agronomiques de Gembloux.
- [36] Carballeira, A., Devesa, C., Retuerto, R., Santillan, E., Uceda, F., 1983. Bioclimatología de Galicia, Xunta de Galicia-Fundación Barrie de la Maza.
- [37] Castroviejo M., 1988. Fitoecología de los montes de Buio y Sierra del Xistral (Lugo), Consellería de Agricultura, Gandería e Pesca, Xunta de Galicia.
- [38] Aussenac G., 2000. Interactions between forest stands and microclimate: Ecophysiological aspects and consequences for silviculture, *Ann. Sci. For.* 57 (3): 287-301. [28] Guilley E., Herve J.C., Nepveu G., 2004. The influence of site quality, silviculture and region on wood density mixed model in *Quercus petraea* Liebl., *For. Ecol. Manag.* 189 (1-3): 111-121.
- [39] Retuerto R., Carballeira A., 1990. Phytoecological importance, mutual redundancy and phytological threshold values of certain climatic factors, *Vegetatio* 90: 47-62.
- [40] Johnson P.S., Shifley S.R., Rogers R., 2002. The ecology and silviculture of oaks, CABI Publishing.
- [42] Thornthwaite C.W., Mather J., 1955. The water balance, *Climatology* 8: 1-104.
- [43] Gaussen H., 1955. Détermination des climats par la méthode des courbes ombrothermiques, *Compt. Rend. Hebd. Séances Acad. Sci.* 240: 642-644.
- [44] Bravo-Oviedo A., Montero G., 2005. Site index in relation to edaphic variables in stone pine (*Pinus pinea* L.) stands in Southwest Spain, *Ann. Sci. For.* 62 (1): 61-72.
- [45] Russell J.S., Moore A.W., 1968. Comparison of different depth weightings in the numerical analysis of anisotropic soil profile data, *Proc. 9th. Int. Cong. Soil Sci.*, 4: 205-213.
- [46] Pardé J., Bouchon J., 1988. Dendrométrie, E.N.G.R.E.F., 2ª ed., Nancy.
- [47] Claessens H., Pauwels D., Thibaut A., Rondeux J., 1999. Site index curves and autoecology of ash, sycamore and cherry in Wallonia (Southern Belgium), *Forestry* 72 (3): 171-182.
- [48] Assmann E., 1970. The principles of forest yield study, Pergamon Press, Oxford, New York.
- [50] Walpole R.E., Myers R.H., Myers S.L., 1999. Probabilidad y estadística para ingenieros, 6ª ed., Prentice Hall, Londres.
- [51] Gaines S.D., Denny M.W., 1993. The largest, smallest, highest, lowest, longest and shortest: extremes in ecology, *Ecology* 74: 1677-1692.
- [52] Daget Ph., Godron M., 1982. Analyse fréquentielle de l'écologie des espèces dans les communautés, Masson.
- [53] Aramburu M.P., Escribano R., Martínez E., Sáenz D., 1984. Análisis de la distribución de

- Quercus pyrenaica* Willd en el Sistema Central, E.T.S.I.M., Madrid. [49] Hart H.M.F., 1928. Stamtal en dunning; een orienteerend onderzoek naar de beste plantwijdte en duningswijze loor den djati, Veenman & Zonen.
- [54] Hill M.O., Bunce R.G.H., Shaw M.W., 1975. Indicator species analysis: a divisive polythetic method of classification, and its application to a survey of native pinewoodws in Scotland, *J. Ecol.* 63: 597-613.
- [55] Hix D.M., 1988. Multifactor classification and analysis of upland hardwood forest ecosystems of the Kichapoo River watershed, southwestern Wisconsin, *Can. J. For. Res.* 18: 1405-1415.
- [56] Martínez E., Ayuga E., González C., 1992. Estudio comparativo de distintas funciones núcleo para la obtención del mejor ajuste según el tipo de datos, *Qüestió* 16: 3-26.
- [57] Hill M.O., 1979a. Decorana: A fortran program for detrended correspondence analysis and reciprocal averaging, Cornell Univ., New York.
- [58] Hill M.O., 1979b. Twinspan: A fortran program for arranging multivariate data in an ordered two-way table by classification of the individuals and attributes, Cornell Univ., New York.
- [59] ter Braak C.J.F., 1994. Canonical community ordination. Part I: Basic theory and linear methods, *Ecoscience* 1: 127-140.
- [60] Kent M., Coker P., 1996. Vegetation description and analysis. A practical approach. John Wiley & Sons, New York.
- [61] Pisces Conservation L.T.D., 1999. Community Analysis Package versión 1.42. A program to search for structure in ecological community data, England.
- [62] Ryan T.P., 1997. Modern regression methods, John Wiley & Sons.
- [63] Sas Institute Inc., 2004. SAS/STAT® 9.1. User's Guide. Cary: SAS Institute Inc.
- [64] Wilde S.A., 1946. Forest soils and forest growth, *Chronica Botanica Comp.*
- [65] Gandullo J.M., 1974. Ensayo de evaluación cuantitativa de la insolación en función de la orientación y de la pendiente del terreno, *Anales INIA, Serie Recursos Naturales* 1: 95-107. [41] Thornthwaite C.W., Mather J., 1957. Instructions and Tables for Computing Potential Evapotranspiration and the Water Balances, Centerton, New Jersey.
- [66] Andre F., Ponette Q., 2003. Comparison of biomass and nutrient content between oak (*Quercus petraea*) and hornbeam (*Carpinus betulus*) trees in a coppice-with-standards stand in Chimay (Belgium), *Ann. Sci. For.* 60 (6): 489-502.
- [67] Morh D., Topp W., 2005. Hazel improves soil quality of sloping oak stands in a German low mountain range, *Ann. Sci. For.* 62(1): 23-30.

ANNEX: List of parameters studied

(1) Mean altitude (ALT), (2) Mean slope (SLP), (3) Orientation (ORI), (4) Sunshine (SUN), (5) Soil depth (DPTH), (6) Distance from sea (DS), (7) Annual precipitation (AP), (8) Winter precipitation (WP), (9) Spring precipitation (SpP), (10) Summer precipitation (SuP), (11) Autumn precipitation (AP), (12) Mean annual temperature (MT), (13) Mean temperature in the hottest month (HMT), (14) Mean temperature in the coldest month (CMT), (15) Annual mean of absolute maximum temperatures (AMT), (16) Annual mean of absolute minimum temperatures (AmT), (17) Temperature range (TR), (18) Annual evapotranspiration (AE), (19) Temperature index (TI), (20) Sum of surpluses (SUR), (21) Sum of deficits (DEF), (22) Hydric index (HI), (23) Duration of drought (DD), (24) Total pH in H₂O (pH), (25) Surface pH in H₂O (spH), (26) Total organic matter (OM), (27) Surface organic matter (sOM), (28) Total nitrogen (N), (29) Surface nitrogen (sN), (30) Total C/N ratio (C/N), (31) Ratio of surface carbon/nitrogen (sC/N), (32) total available phosphorous (P), (33) Surface available phosphorous (sP), (34) Total exchangeable

potassium (K), (35) Surface exchangeable potassium (sK), (36) Total exchangeable calcium (Ca), (37) Surface exchangeable calcium (sCa), (38) Total exchangeable magnesium (Mg), (39) Surface exchangeable magnesium (sMg), (40) Percentage of fine earth (FE), (41) Percentage of total gravel (GR), (42) Number of trees per hectare (NT/ha), (43) Number of non inventoriable *Quercus petraea* specimens (NNQ), (44) Number of dead specimens of *Quercus petraea* (NDQ), (45) Total number of non-inventoriable trees (TNNT), (46) Total number of dead trees (TNNT), (47) Regenerated saplings (REG), (48) Basimetric area per hectare (BA), (49) Mean arithmetic diameter (MAD), (50) Mean quadratic diameter (MQD), (51) Standard deviation of the diameter distribution (SDD), (52) Coefficient of variation of the diameter distribution (CVD), (53) Dominant diameter (DOD), (54) Mean arithmetic height (MAH), (55) Mean quadratic height (MQH), (56) Standard deviation of the height distribution (DEH), (57) Coefficient of variation of the height distribution (CVH), (58) Assmann's dominant height (HDA), (59) Hart's index (HAI) and (60) Czarnowski's index (CZI).

Table I. Sample plots descriptive data.

Number	Name	Province	Surface area (m ²)	Altitude (m)	Slope (%)	Orientation
1	Ancares 1	Lugo	1200	1220	52	North
2	Ancares 2	Lugo	600	1000	57	North
3	Ancares 3	Lugo	600	965	81	East
4	Ancares 4	Lugo	1200	1310	53	East
5	Ancares 5	Lugo	875	1215	50	West
6	Ancares 6	Lugo	1000	1190	53	West
7	Baleira	Lugo	1050	840	42	North
8	Candín 1	León	600	1230	39	North
9	Candín 2	León	400	1070	63	North
10	Cangas 1	Asturias	500	1000	75	East
11	Cangas 2	Asturias	500	1100	75	East
12	Cerredo 1	Asturias	500	1120	36	North
13	Cerredo 2	Asturias	400	975	27	North
14	Cerredo 3	Asturias	1200	1040	23	West
15	Cerredo 4	Asturias	1000	1165	78	East
16	Cerredo 5	Asturias	900	1120	37	North
17	Cerredo 6	Asturias	1200	1080	43	North
18	Cortes	Asturias	1200	860	28	East
19	Courel 1	Lugo	480	1165	70	North
20	Courel 2	Lugo	600	1395	49	North
21	Courel 3	Lugo	750	1000	32	North
22	Fondos 1	Asturias	400	1110	74	North
23	Fondos 2	Asturias	400	960	63	West
24	Ibías 1	Asturias	600	900	67	North
25	Ibías 2	Asturias	600	800	54	North
26	Ibías 3	Asturias	750	750	41	East
27	Lena 1	Asturias	750	970	28	East
28	Lena 2	Asturias	750	1075	59	South
29	Meira	Lugo	600	665	25	East
30	Palacios 1	León	800	1160	73	East
31	Palacios 2	León	875	1035	47	North
32	Palacios 3	León	875	1180	57	East
33	Palacios 4	León	500	1145	35	East
34	Palacios 5	León	900	1025	90	East
35	Palacios 6	León	700	1000	26	North
36	Pastoriza	Lugo	600	540	8	North
37	Pintinidoira 1	Lugo	500	1200	58	North
38	Pintinidoira 2	Lugo	400	1050	52	North
39	Pontenova 1	Lugo	800	600	53	North
40	Pontenova 2	Lugo	1000	550	58	West
41	Pontenova 3	Lugo	800	690	47	West
42	Suarbol 1	León	600	1175	42	East
43	Suarbol 2	León	875	1165	38	North
44	Suarbol 3	León	600	1245	51	North
45	Teverga 1	Asturias	500	995	42	West
46	Teverga 2	Asturias	900	1030	30	South
47	Teverga 3	Asturias	750	1160	24	West
48	Teverga 4	Asturias	750	1125	21	West
49	Villablino 1	León	600	1235	37	North
50	Villablino 2	León	500	1230	51	West
51	Villablino 3	León	400	1375	57	East
52	Villablino 4	León	500	1400	45	West

Table II. Descriptive statistics of the physiographic and climatic parameters (n=52) (SD: standard deviation, CV: coefficient of variation). The variables without units are adimensional.

Parameter	Mean	SD	CV (%)	Maximum	Minimum
ALT (m)	1053.3	196.3	18.6	1395	540
SLP (%)	48.4	17.6	36.4	90	7.9
ORI (°)	165.3	122.4	74.0	359.0	0.0
SUN	0.8	0.3	37.5	1.3	0.2
DPTH (cm)	103.6	44.6	43.1	190.0	25.0
DS (Km)	84.8	20.8	24.5	129.0	34.0
AP (mm)	1589.5	249.1	15.7	2006.0	1150.0
WP (mm)	495.1	150.2	30.3	711.0	310.2
SpP (mm)	397.0	76.3	19.2	542.8	246.1
SuP (mm)	192.5	28.5	14.8	227.2	143.5
AP (mm)	481.0	76.7	15.9	582.2	334.5
MT (°C)	8.8	2.0	22.7	11.0	5.0
HMT (°C)	15.9	1.9	11.9	17.4	12.3
CMT (°C)	2.5	2.1	84.0	5.4	-0.8
AMT (°C)	20.8	2.8	13.5	23.4	17.7
AmT (°C)	-2.6	2.9	111.5	-0.1	-5.9
TR (°C)	13.3	0.7	5.3	14.8	12.0
AE (mm)	596.2	52.6	8.8	654	497
TI	194.8	42.0	21.6	240.0	101.0
SUR (mm)	1228.5	160.6	13.1	1486.0	763.0
HI	246.2	24.0	9.7	273.0	224.0
DD (n° months)	0.1	0.2	200.0	1.0	0.0

See list of parameters in the annex. The parameter DEF was not analyzed because of the non-existence of hydric deficit in most of the plots.

Table III. Descriptive statistics of the edaphic parameters (n=52) (SD: standard deviation, CV: coefficient of variation). Variables without units are adimensional.

Parameter	Mean	SD	CV (%)	Maximum	Minimum
pH	4.73	0.36	7.6	5.65	4.23
sPH	4.59	0.43	9.4	5.60	3.90
OM (%)	7.83	4.04	51.6	19.83	1.82
sOM (%)	10.09	4.56	45.2	24.31	2.89
N (%)	0.25	0.11	44.0	0.55	0.07
sN (%)	0.32	0.13	40.6	0.72	0.10
C/N	17.92	3.75	20.9	23.14	8.83
sC/N	14.07	4.06	28.9	20.99	6.63
P (ppm)	9.53	7.49	78.6	24.80	0.93
sP (ppm)	12.57	14.49	115.3	52.69	0.76
K (ppm)	90.04	52.90	58.8	275.09	25.69
sK (ppm)	114.68	56.19	49.0	264.51	37.10
Ca (ppm)	203.21	275.67	135.7	1135.07	13.19
sCa (ppm)	291.29	342.47	117.6	1431.85	14.01
Mg (ppm)	45.54	42.50	93.3	164.96	3.76
sMg (ppm)	61.88	46.78	75.6	198.87	4.79
FE (%)	43.64	11.45	26.2	67.20	17.80
GR (%)	56.36	11.45	20.3	82.20	32.80

See list of parameters in the annex

Table IV. Descriptive statistics of the dendrometric/silvicultural parameters (n=52) (SD: standard deviation, CV: coefficient of variation). Variable without units are adimensional.

Parameter	Mean	SD	CV (%)	Maximum	Minimum
NT/ha (No ha ⁻¹)	990.0	495.9	50.1	2950.0	267.0
NNQ	9.3	18.6	200.0	78.0	0.0
NDQ	10.3	10.4	101.0	47.0	0.0
TNNT	57.0	72.4	127.0	348.0	0.0
TNDT	11.9	11.7	98.3	47.0	0.0
REG	2.4	3.0	125.0	10.0	0.0
BA (m ² ha ⁻¹)	31.3	6.8	21.7	54.0	17.6
MAD (cm.)	21.4	7.9	36.9	41.5	2.4
MQD(cm)	23.8	7.2	30.3	43.2	12.0
SDD (cm)	8.5	2.5	29.4	15.5	3.7
CVD (%)	39.7	11.2	28.2	65.0	16.0
DOD (cm)	35.2	7.6	21.6	55.1	18.7
MAH (m)	15.7	3.2	20.4	23.9	11.3
MQH (m)	4.0	0.4	10.0	4.9	3.4
DEH (m)	3.0	0.9	30.0	5.8	1.1
CVH (%)	15.8	6.3	39.9	39.0	6.5
ADH (m)	17.5	3.9	22.3	27.3	10.2
HAI (%)	20.5	5.9	28.8	37.0	11.0
CZI	23.1	11.1	48.1	56.2	7.0

See list of parameters in the annex

Table V. Final groups resulting from application of the Twinspan programme.

Group	Discriminant parameters	Plots
A	ALT > M – DE	11, 20, 21
B	ALT < M – DE	7, 29, 36, 39, 40, 41
C	M – DE < WP < M + DE	2, 3
D	WP < M – DE	18, 27, 28, 45, 46, 47, 48
E	spH < M – DE	5, 6, 42, 43, 44
F	M – DE < spH < M + DE	1, 4, 8, 9
G	sK < M – DE	12, 13, 14, 15, 26
H	sK > M + DE	22, 23, 24, 25
I	sOM > M + DE	10, 11, 16, 17
J	TI < M – DE	37, 38
K	M – DE < Ca < M + DE	30, 31, 32, 33, 49, 50, 52
L	Ca > M + DE	34, 35, 51

Table VI. Coefficients of Pearson's linear correlation between the dendrometric/silvicultural parameters.

	NT/h	NNQ	NDQ	TNNT	TNDT	REG	BA	MAD	MQD	SDD	CVD	DOD	MAH	MQH	DEH	CVH	ADH	HAI	CZI
NT/h	1,000	,392**	,596**	n.s.	,511**	n.s.	n.s.	-.604**	-.782**	-.510**	n.s.	-.602**	-.455**	-.387**	n.s.	,372**	-.365**	-.491**	,584**
NNQ		1,000	n.s.	,281*	n.s.	n.s.	n.s.	-.294*	n.s.	n.s.	n.s.	n.s.	-.427**	-.361**	n.s.	n.s.	-.534**	,348*	n.s.
NDQ			1,000	n.s.	,879**	n.s.	n.s.	-.357**	-.444**	n.s.	n.s.	-.324*	n.s.	n.s.	n.s.	,312*	n.s.	-.336*	,399**
TNNT				1,000	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
TNDT					1,000	n.s.	n.s.	-.372**	-.454**	n.s.	,274*	-.350*	n.s.	n.s.	n.s.	,329*	n.s.	-.324*	,339*
REG						1,000	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
BA							1,000	,373**	,379**	,303*	n.s.	,398**	,504**	,472**	n.s.	n.s.	,435**	n.s.	,367**
MAD								1,000	,823**	-.604**	-.305*	,660**	,604**	,547**	n.s.	-.351*	,472**	n.s.	n.s.
MQD									1,000	,596**	n.s.	,784**	,639**	,554**	n.s.	-.350*	,508**	,361**	-.329*
SDD										1,000	,598**	,731**	,400**	,337*	n.s.	n.s.	,433**	n.s.	n.s.
CVD											1,000	n.s.	n.s.	n.s.	n.s.	,667**	n.s.	n.s.	n.s.
DOD												1,000	,493**	,429**	n.s.	n.s.	,465**	n.s.	n.s.
MAH													1,000	,774**	n.s.	-.353*	,885**	-.285*	,383**
MQH														1,000	n.s.	-.305*	,680**	n.s.	n.s.
DEH															1,000	n.s.	,312*	-.291*	n.s.
CVH																1,000	n.s.	n.s.	n.s.
ADH																	1,000	-.510**	,413**
HAI																		1,00	-.781**
CZI																			1,000

Levels of significance (s): *, s > 95%; **, s > 99%; n.s., non significant

Figure 1. Location of the sampling plots in the study area within the Iberian Peninsula.



Figure 2. Central and marginal physiographic-climatic habitats of *Quercus petraea* in the northwest Iberian Peninsula.




Figure 3. Central and marginal edaphic habitats of *Quercus petraea* in the northwest Iberian Peninsula.




Figure 4. Spatial distribution of the plots identified according to the groups obtained by discriminant analysis.

